

THE INTERNATIONAL SERIES OF
MONOGRAPHS ON CHEMISTRY

1. J. D. Lambert: *Vibrational and rotational relaxation in gases*
2. N. G. Parsonage and L. A. K. Staveley: *Disorder in crystals*
3. G. C. Maitland, M. Rigby, E. B. Smith, and W. A. Wakeham:
Intermolecular forces: their origin and determination
4. W. G. Richards, H. P. Trivedi, and D. L. Cooper: *Spin-orbit coupling
in molecules*
5. C. F. Cullis and M. M. Hirschler: *The combustion of organic polymers*
6. R. T. Bailey, A. M. North, and R. A. Pethrick: *Molecular motion in
high polymers*
7. Atta-ur-Rahman and A. Basha: *Biosynthesis of indole alkaloids*
8. J. S. Rowlinson and B. Widom: *Molecular theory of capillarity*
9. C. G. Gray and K. E. Gubbins: *Theory of molecular fluids. Volume 1:
Fundamentals*
10. C. G. Gray and K. E. Gubbins: *Theory of molecular fluids. Volume 2:
Applications (in preparation)*
11. S. Wilson: *Electron correlation in molecules*
12. E. Haslam: *Metabolites and metabolism*
13. G. R. Fleming: *Chemical applications of ultrafast spectroscopy*
14. R. R. Ernst, G. Bodenhausen, and A. Wokaun: *Principles of nuclear
magnetic resonance in one and two dimensions*
15. M. Goldman: *Quantum description of high-resolution NMR in liquids*
16. R. G. Parr and W. Yang: *Density-functional theory of atoms and
molecules*
17. J. C. Vickerman, A. Brown, and N. M. Reed (editors): *Secondary ion
mass spectrometry: principles and applications*
18. F. R. McCourt, J. Beenakker, W. E. Köhler, and I. Kušćer:
Nonequilibrium phenomena in polyatomic gases, Volume 1: Dilute gases
19. F. R. McCourt, J. Beenakker, W. E. Köhler, and I. Kušćer:
*Nonequilibrium phenomena in polyatomic gases, Volume 2: cross
sections, scattering, and rarefied gases*
20. T. Mukaiyama: *Challenges in synthetic organic chemistry*
21. P. Gray and S. K. Scott: *Chemical oscillations and instabilities: non-
linear chemical kinetics*
22. R. F. W. Bader: *Atoms in molecules: a quantum theory*
23. J. H. Jones: *The chemical synthesis of peptides*
24. S. K. Scott: *Chemical chaos*
25. M. S. Child: *Semiclassical mechanics with molecular applications*
26. D. T. Sawyer: *Oxygen chemistry*

The Chemical Synthesis of Peptides

John Jones

Balliol College, Oxford

Best Available Copy

CLARENDON PRESS · OXFORD
1991

Oxford University Press, Walton Street, Oxford OX2 6DP

Oxford New York Toronto
Delhi Bombay Calcutta Madras Karachi
Petaling Jaya Singapore Hong Kong Tokyo
Nairobi Dar es Salaam Cape Town
Melbourne Auckland
and associated companies in
Berlin Ibadan

Oxford is a trade mark of Oxford University Press

Published in the United States
by Oxford University Press, New York

© John Jones, 1991

All rights reserved. No part of this publication may be reproduced,
stored in a retrieval system, or transmitted, in any form or by any means,
electronic, mechanical, photocopying, recording, or otherwise, without
the prior permission of Oxford University Press

British Library Cataloguing in Publication Data

A catalogue record for this book is
available from the British Library

Library of Congress Cataloging in Publication Data

Jones, John Henry, 1942 Apr. 29–
The chemical synthesis of peptides / John Jones.
(International series of monographs on chemistry; 23)
Includes index.

1. Peptides—Synthesis. I. Title. II. Series.
QP552.P4J66 1991 574.19'296—dc20 90-23913

Printed in Great Britain by
Bookcraft (Bath) Ltd
Midsomer Norton, Avon

ACKNOWLEDGEMENTS


Dr Geoffrey Young first interested me in peptide chemistry more than a quarter of a century ago, and sustained my enthusiasm for it over many years. I am glad to have an opportunity to acknowledge my great obligation to him here.

Many correspondents have sent reprints and given assistance in other ways, but I am especially grateful to Professor David Evans, Dr Bruno Kamber, and Dr Barbara Rzesotarska for unpublished material, and to Professors Bruce Merrifield, John Sheehan, and Theodor Wieland for permission to quote remarks made by them.

Mr Ian Eggleston kindly read the whole book in draft and made many helpful comments. Most of it was written during sabbatical leave in the academic year 1989–90. This privilege inevitably meant extra burdens for colleagues, both in Balliol and the Dyson Perrins Laboratory: I appreciate with special warmth the Reverend Dr Douglas Dupree's willingness to be Acting Dean of Balliol for a year, and the patient skill of my secretary, Miss Rachel Johnson.

Balliol College, Oxford
May 1990

J.H.J.




case), and the progressive accumulation of over-reaction products limits the extent to which repetitive approach can be taken without difficult purification problems. The thio-analogues or NTAs (25) have also been used for peptide synthesis.⁶⁵ They are less prone to yield over-reaction products than NCAs, as the thiocarbanic acids produced by their aminolysis are less fragile. They are preferred in the special cases of glycine and histidine (the NCAs of which are especially subject to troublesome side-reactions), but unfortunately they are not secure against racemization.

5.1.2 Racemization*


Consider the synthesis of an all-L peptide comprising n chiral residues, from optically pure α -amino acids. If the operations needed for the incorporation of each residue result in conversion of a small fraction f of each residue to the D-form, and the epimers are carried through without separation, then the end product will consist of the required all-L peptide and a blend of other peptides in approximate proportions $(1-f)^n:fn$. For a synthesis of a 50-residue peptide in which 1% D-residue formation takes place at each stage, only half the final product will have the required all-L stereochemistry. The other 50% will consist mainly of about 1% each of all the 50 possible epimers with one D-residue. This will in general pose a prohibitive purification problem, and racemization in peptide synthesis has therefore been closely studied, with a view to defining the conditions under which it is minimal.⁶⁶⁻⁶⁸

Except for special cases (e.g., synthesis with *N*-methylamino acids: see Section 6.2.1), racemization is an almost exclusively base-induced side-reaction, and in practice is only a matter for serious concern at the activation and coupling stages of a synthesis. There are two important mechanisms.

*This term is used in peptide chemistry in a loose way which not only covers the strict sense as defined in most general organic chemistry texts (conversion of an enantiomer to a mixture of enantiomers), but also embraces partial epimerisation, whereby there is loss of chiral integrity at one out of two or more chiral centres, resulting in the formation of a mixture of epimers (i.e. diastereoisomers differing at one chiral centre).





Scheme 5.30.



Scheme 5.31.

5.1.2.1 Direct enolization


Deprotonation at the α -carbon of an α -amino-acid residue results in racemization, because the carbanion intermediate can reprotonate on either side (Scheme 5.30). This has been called the 'direct exchange' mechanism, an inappropriate expression, because under some circumstances⁶⁹ racemization is much faster than exchange with the proton pool, implying that an ion pair is formed in which the ions are jostled about by solvent molecules and change their relative orientations without being divorced from each other, so that reprotonation can return the original proton to either side of the chiral centre ('isomerization': Scheme 5.31). The rate of racemization by direct enolization depends on the catalysing base, the solvent, and the electron-withdrawing effects of the groups P , R , and X around the chiral centre (26). When $X = \text{NH-}$, O-alkyl , or O- , it is in most cases negligible,



Scheme 5.33.

does not arise because there is no hydrogen at the α -carbon. When the amino-nitrogen of the activated residue is acylated with a simple acyl group (acetyl, benzoyl, etc.), or with a peptide chain, cyclization to the oxazolone occurs easily with most good leaving groups X, and gross or even complete racemization may ensue. But oxazolone formation is not so facile when the acyl substituent is an alkoxycarbonyl protecting group. Indeed, the process was held to be impossible until 1977.⁷² Furthermore, the alkoxyoxazolones **28** are both less easily racemized and more easily aminolysed than are the oxazolones **27** derived from simple acylamino acids. The activation of ordinary Z, Boc and Fmoc amino acids, etc., and their coupling with amino components is consequently not attended by the danger of racemization under normal conditions. This is a pivotal fact on which much of modern peptide synthesis turns. The reason for the contrast between, e.g., Z and benzoyl amino acids has not been fully explained, but a major factor is probably the lower acidity of Bz1OCNH- compared to PhCONH- . In Scheme 5.32, the ring closure is shown as a specific base catalysed process, which has been demonstrated to be so in one set of circumstances.⁷² It might be a concerted general base catalysed process under other conditions, but in either case lowering the acidity of the NH would be expected to diminish the rate of oxazolone formation.


A mechanism for oxazolone formation which does not require base assistance is also possible (Scheme 5.33), but ordinary amino acid derivatives do not cyclize this way except under very vigorous activation. *N*-Methyl- α -amino acid derivatives do, however, and give optically labile oxazolonium cations (see Scheme 6.29) even under normal activation conditions.⁷⁴ base catalysis is impossible because there is no NH for it to operate through, so cyclization can only occur by attack of the neutral amide oxygen on the activated carbonyl, which is easier with $-\text{CONMe-}$ than $-\text{CONH-}$ because




and basic deprotection procedures other than saponification (see Section 4.1.1) are generally completely safe. During activation and coupling, the risk is rather more significant, but the danger is over once coupling is complete. Racemization is fastest with strongly electron-withdrawing groups X (i.e. most good leaving groups), and unhindered strong bases in dipolar aprotic solvents like DMSO and DMF. Inessential exposure to strong bases is clearly to be avoided, but the best way of responding to the other factors is not so obvious, because what matters is not the rate of racemization *per se*, but the balance between the rates of racemization and coupling, and this is much more difficult to make reliable generalizations about. Fortunately, with the exception of (a) a few special amino acids (α -arylglycines present quite a serious problem), and (b) couplings which are inordinately slow, the amount of racemization which actually takes place by this pathway is very slight indeed.

5.1.2.2 The oxazolone mechanism

Activated acylamino acids and peptides cyclize under the influence of base to give oxazolones [**27**; strictly '5(4*H*)-oxazolones', formerly '2-oxazolin-5-ones', archaically 'azlactones']. The oxazolones so formed are themselves activated towards aminolysis, and reaction with amino components leads ultimately to peptides, but since their racemization via stabilized anions is usually fast compared to the rate of peptide bond formation, any peptide thus produced is largely racemized (Scheme 5.32). Oxazolones are actually useful (e.g., references 70 and 71; see Scheme 6.31) for the activation of α -dialkyl- α -amino acid residues, where the question of racemization by base



Scheme 5.32. Conditions: basic.



of electron release by the methyl group. Conformational restraints fortunately prevent this happening with proline derivatives, under all except the most extreme conditions.

Whether or not oxazolone-mediated racemization accompanies the activation and coupling of susceptible protected peptide acids depends on the leaving group. It does not seem to do so to a significant extent with acyl azide intermediates, in model systems at least, perhaps because these owe their aminolytic reactivity to intramolecular general base catalysis (see Section 5.1.1.2), to which oxazolone formation is indifferent because the nucleophile bears no hydrogen. And the risk is also small with the DCCI-HOBt procedure (see Section 5.1.1.4), because HOBt rapidly intercepts the activated species which might otherwise degenerate into oxazolones. The reactive HOBt ester intermediate favours aminolysis over oxazolone formation, possibly in part for the same reason as suggested for azides.

With activated protected peptides, the direct enolization and oxazolone mechanisms both provide pathways for the racemization of the carboxy-terminal residue. The formation of an oxazolone also threatens the chiral integrity of the penultimate residue, because the carbanion **29** is stabilised. Racemization at that residue has been observed at moderate levels in model experiments (e.g., reference 75), but has not so far been recognized as a real problem in actual syntheses, except when deliberate oxazolone formation is used to drive the coupling of carboxy components terminating in -XaaAibOH.⁷¹


5.2 The use of enzymes

This book is concerned with the chemical synthesis of peptides, so the inclusion of enzymic methods may raise a few eyebrows, but the use of enzymes as reagents in preparative organic chemistry,⁷⁶ without special homage to their biological origin, is burgeoning. A few remarks on enzymatic peptide bond formation^{77,78} therefore seem called for.


Nature provides a wide range of proteolytic enzymes which can in principle be perverted to catalyse peptide bond formation by manipulating the conditions. There are two strategies for doing this. The first is dependent on thermodynamic control, the equilibrium in Scheme 5.34 (which



Scheme 5.34.



Scheme 5.35.



Scheme 5.36. Conditions: i, 2 equiv. HPhOMe/pH7/thermolysin (the dipeptide salt precipitates); ii, HCl, then catalytic transfer hydrogenolysis with $HCO_2NH_4/Pd(C)/MeOH$.

favours hydrolysis overwhelmingly under normal conditions) being somewhat displaced in favour of peptide bond formation. This can be achieved by employing protecting groups which will ensure precipitation of the peptide, or by using biphasic systems so that the peptide passes out of the aqueous phase into an organic solvent as it is formed, or by using water-miscible organic solvents which perturb the dissociation constants of the components and shift the balance of the equilibrium. The second strategy exerts kinetic control by arranging for an amino component nucleophile to compete with water for an acyl-enzyme intermediate (Scheme 5.35). The advantages of an enzymatic synthesis are the mild conditions, freedom from racemization and the need for side-chain protection, the possibility of using immobilized enzyme technology^{79,80} with catalyst recovery, and the scope for industrial scale-up. Many examples have been reported. The synthesis⁸¹ of the synthetic sweetener aspartame (**30**) is one of particular interest which has been developed for commercial application, and is also simple enough to be an undergraduate exercise⁸² (Scheme 5.36). There are disadvantages, however. With peptides longer than dipeptides, there is the danger that while the protease is being persuaded to work backwards in creating a peptide bond at one point, it will remember the purpose for which evolution devised it and dismantle another somewhere else. No new case can be treated as

36. Sheehan, J.C. (1982). *The enchanted ring*. MIT Press, Cambridge, Massachusetts.
37. Sheehan, J.C. and Hess, G.P. (1955). *Journal of the American Chemical Society*, **77**, 1067-1068.
38. Weygand, F., Hoffmann, D., and Wünsch, E. (1966). *Zeitschrift für Naturforschung*, **21B**, 426-428.
39. Wünsch, E. and Drees, F. (1966). *Chemische Berichte*, **99**, 110-120.
40. König, W. and Geiger, R. (1970). *Chemische Berichte*, **103**, 788-798.
41. Balcom, B.J. and Petersen, N.O. (1989). *Journal of Organic Chemistry*, **54**, 1922-1927.
42. Sarantakis, D., Teichman, J., Lien, E.L., and Fenichel, R.L. (1976). *Biochemical and Biophysical Research Communications*, **73**, 336-342.
43. Sheehan, J.C., Cruickshank, P.A., and Boshart, G.L. (1961). *Journal of Organic Chemistry*, **26**, 2525-2528.
44. Klausner, Y.S. and Bodanszky, M. (1972). Coupling reagents in peptide synthesis. *Synthesis*, 453-463.
45. Woodward, R.B., Olofson, R.A., and Mayer, H. (1966). *Tetrahedron*, Supplement 8, Part 1, 321-346.
46. Kemp, D.S., Wrobel, S.J., Wang, S.-W., Bernstein, Z., and Rebek, J., Jr. (1974). *Tetrahedron*, **30**, 3969-3980.
47. Mukaiyama, T., Matsueda, R., and Ueki, M. (1979). The oxidation-reduction condensation. In *The peptides*, Vol. 2 (ed. E. Gross and J. Meienhofer), Chapter 5. Academic Press, London.
48. Le Nguyen, D. and Castro, B. (1988). In *Peptide chemistry 1987* (ed. T. Shiba and S. Sakakibara), pp 231-238. Protein Research Foundation, Osaka.
49. Steinauer, R., Chen, F.M.F., and Benoiton, N.L. (1989). *International Journal of Peptide and Protein Research*, **34**, 295-298.
50. (a) Coste, J., Dufour, M.-N., Le-Nguyen, D., and Castro, B. (1990). In *Peptides: chemistry structure and biology* (ed. J.E. Rivier and G.R. Marshall), pp 885-888. ESCOM Science Publishers, Leiden.
(b) Coste, J., Le-Nguyen, D., and Castro, B. (1990). *Tetrahedron Letters*, **31**, 205-208.
51. Knorr, R., Trzeciak, A., Bannwarth, W., and Gillesen, D. (1989). *Tetrahedron Letters*, **30**, 1927-1930. Much work on phosphonium and uronium coupling reagents appeared while this book was in the final stages of preparation: for a useful bibliography see *Novabiochem Letters*, number 1/90 [pub. Novabiochem (UK) Ltd].
52. Menger, F.M. and Smith, J.H. (1972). *Journal of the American Chemical Society*, **94**, 3824-3829.

53. Bodanszky, M. (1979). Active esters in peptide synthesis. In *The peptides*, Vol. 1 (ed. E. Gross and J. Meienhofer), Chapter 3. Academic Press, London.
54. Atherton, E. and Sheppard, R.C. (1989). In *Solid phase peptide synthesis: a practical approach*, p 78. IRL Press at Oxford University Press, Oxford.
55. Jaouadi, M., Martinex, J., and Castro, B. (1987). *Journal of Organic Chemistry*, **52**, 2364-2367.
56. Kisfaludy, L. and Nyéki, O. (1975). *Acta Chimica Academiae Scientiarum Hungaricae*, **88**, 343-345.
57. Bell, J.R., Jones, J.H., and Webb, T.C. (1975). *International Journal of Peptide and Protein Research*, **7**, 235-242.
58. Fairweather, R. and Jones, J.H. (1972). *Journal of the Chemical Society Perkin Transactions I*, 1908-1915.
59. Jones, J.H. and Young, G.T. (1968). *Journal of the Chemical Society, Section C*, 436-441.
60. Kricheldorf, H.R. (1987). *α -Aminoacid-N-carboxyanhydrides and related heterocycles*. Springer-Verlag, Berlin.
61. Blacklock, T.J., Hirschmann, R., and Veber, D.F. (1987). The preparation and use of N-carboxyanhydrides and N-thiocarboxyanhydrides for peptide bond formation. In *The peptides*, Vol. 9 (ed. S. Udenfreind and J. Meienhofer), Chapter 2. Academic Press, London.
62. Fasman, G.D. (1967). *Poly- α -amino acids*. Edward Arnold, London.
63. Blacklock, T.J., Shuman, R.F., Butcher, J.W., Shearin, W.E., Jr., Budavari, J., and Grenda, V.J. (1988). *Journal of Organic Chemistry*, **53**, 836-844.
64. Denkewalter, R.G., Schwamm, H., Strachan, R.G., Beesley, T.E., Veber, D.F., Schoenewaldt, E.F., Barkemeyer, H., Palevda, W.J., Jr., Jacob, T.A., and Hirschmann, R. (1966). *Journal of the American Chemical Society*, **88**, 3163-3164.
65. Dewey, R.S., Schoenewaldt, E.F., Joshua, H., Palevda, W.J., Jr., Schwamm, H., Barkemeyer, H., Arison, B.H., Veber, D.F., Strachan, R.G., Milkowski, J., Denkewalter, R.G., and Hirschmann, R. (1971). *Journal of Organic Chemistry*, **36**, 49-59.
66. Kemp, D.S. (1979). Racemization in peptide synthesis. In *The peptides*, Vol. 1 (ed. E. Gross and J. Meienhofer), Chapter 7. Academic Press, London.
67. Kovács, J. (1980). Racemization and coupling rates of N $^{\alpha}$ -protected amino acid and peptide active esters: predictive potential. In *The peptides*, Vol. 2 (ed. E. Gross and J. Meienhofer) Chapter 8. Academic Press, London.
68. Benoiton, N.L. (1983). Quantitation and sequence dependence of racemization in peptide synthesis. In *The peptides*, Vol. 5 (ed. E.

- Gross and J. Meienhofer), Chapter 4. Academic Press, London.
69. Kovacs, J., Cortegiano, H., Cover, R.E., and Mayers, G.L. (1971). *Journal of the American Chemical Society*, **93**, 1541-1543.
 70. Leplawy, M.T., Jones, D.S., Kenner, G.W., and Sheppard, R.C. (1960). *Tetrahedron*, **11**, 39-51.
 71. Brückner, H. (1989). In *Chemistry of peptides and proteins* (ed. W.A. König and W. Voelter), pp 79-86. Attempto Verlag, Tübingen.
 72. Jones, J.H. and Witty, M.J. (1977). *Journal of the Chemical Society Chemical Communications*, 281-282.
 73. Kemp, D.S. and Chien, S.W. (1967). *Journal of the American Chemical Society*, **89**, 2745-2746.
 74. McDermott, J.R. and Benoiton, N.L. (1973). *Canadian Journal of Chemistry*, **51**, 2562-2570.
 75. Weygand, F., Prox, A., and König, W. (1966). *Chemische Berichte*, **99**, 1446-1450.
 76. (a) Davies, H.G., Green, R.H., Kelly, D.R., and Roberts, S.M. (1989). *Biotransformations in preparative organic chemistry*. Academic Press, London. (b) Turner, N.J. (1989). Recent advances in the use of enzyme-catalysed reactions in organic synthesis. *Natural Product Reports*, **6**, 625-644.
 77. Jakubke, H.-D. (1987). Enzymatic peptide synthesis. In *The peptides*, Vol. 9 (ed. S. Udenfriend and J. Meienhofer). Academic Press, London.
 78. Kullmann, W. (1987). *Enzymatic peptide synthesis*. CRC Press Inc., Boca Raton, Florida.
 79. Jakubke, H.-D. and Könecke, A. (1987). Peptide synthesis using immobilised proteases. *Methods in Enzymology*, **136**, 178-188.
 80. Okahata, Y. and Iijiro, K. (1988). *Journal of the Chemical Society, Perkin Transactions II*, 91-96.
 81. Isowa, Y., Ohmori, M., Ichikawa, T., Mori, K., Nonaka, Y., Kihara, K., and Oyama, K. (1979). *Tetrahedron Letters*, **28**, 2611-2612.
 82. Lindeberg, G. (1987). In *Peptides 1986* (ed. D. Theodoropoulos), pp 199-202. Walter de Gruyter, Berlin.
 83. Elmore, D.T. (1987). Personal communication, November 1987.
 84. Schultz, P.G. (1989). Catalytic antibodies. *Accounts of Chemical Research*, **22**, 287-294, and *Angewandte Chemie International Edition in English*, **28**, 1283-1295.
 85. Brenner, M. (1966). Critical evaluation of coupling methods. In *Peptides* (ed. H.C. Beyerman, A. van de Linde, and W. Maassen van den Brink), pp 1-7. North-Holland Publishing Co., Amsterdam.
 86. Kemp, D.S. and Vellaccio, F., Jr. (1975). *Journal of Organic Chemistry*, **40**, 3003-3004.
 87. Fotouhi, N., Galakatos, N.G., and Kemp, D.S. (1989). *Journal of*

- Organic Chemistry*, **54**, 2803-2817.
88. Takasaki, M. and Harada, K. (1987). *Journal of the Chemical Society Chemical Communications*, 571-573.
 89. Ojima, I. and Chen, H.-J.C. (1987). *Journal of the Chemical Society Chemical Communications*, 625-626.
 90. Lipshutz, B.H. and Reuter, D.C. (1988). *Tetrahedron Letters*, **29**, 6067-6070.
 91. Sahebi, M., Wipf, P., and Heimgartner, H. (1989). *Tetrahedron*, **45**, 2999-3010.
 92. Duhamel, P., Goument, B., and Plaquevent, J.-C. (1987). *Tetrahedron Letters*, **28**, 2595-2596.
 93. Ugi, I. (1980). The four component synthesis. In *The peptides*, Vol. 2 (ed. E. Gross and J. Meienhofer), Chapter 4. Academic Press, London.
 94. Waki, M. and Meienhofer, J. (1977). *Journal of the American Chemical Society*, **99**, 6075-6082.
 95. Urban, R., Eberle, G., Marquarding, D., Rehn, D., Rehn, H., and Ugi, I. (1976). *Angewandte Chemie International Edition in English*, **15**, 627-628.
 96. Bowers, M.M., Carroll, P., and Joullié, M.M. (1989). *Journal of the Chemical Society Perkin Transactions I*, 857-865.
 97. Maia, H.L., Ridge, B., and Rydon, H.N. (1973). *Journal of the Chemical Society Perkin Transactions I*, 98-105.
 98. Leuchs, H. and Suzuki, U. (1904). *Berichte der Deutschen Chemischen Gesellschaft*, **37**, 3306-3316.
 99. Erdik, E. and Ay, M. (1989). Electrophilic amination of carbanions. *Chemical Reviews*, **89**, 1947-1980.
 100. Evans, D.A., Weber, A.E., Britton, T.C., Ellman, J.A., and Sjogren, E.B. (1987). Asymmetric synthesis of amino acids. In *Peptides* (ed. G.R. Marshall), pp 143-148. ESCOM Science Publishers, Leiden.
 101. Evans, D.A., Britton, T.C., Ellman, J.A., and Dorow, R.L., (1990). *Journal of the American Chemical Society*, **112**, 4011-4030.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.